



Your physician has recommended the following products to aid in your recovery following surgery.

DONJOY® ICEMAN® CLEAR³



THE
CLEAR
ADVANTAGE

WHAT IS IT?

The DonJoy® IceMan® CLEAR³ cold therapy unit helps reduce pain and swelling, speeding up rehabilitation. It provides extended cold therapy for a variety of indications and protocols as directed by a medical professional and utilizes DonJoy's patented semi-closed loop recirculation system, which maintains more consistent and accurate temperatures than other cold therapy units, in an easy to use preset configuration.

WHY USE IT?

It takes the place of ice packs saving you multiple trips to the freezer and does not "weigh down" on your shoulder like an ice pack does. Your physician will give you a prescription with all the necessary details on how long and how often the device should be used for your situation.

DO NOT use this device without a prescription from a physician. Rx only. Your prescription must state a temperature, how long and how often the device should be used and the length of breaks between uses. DO NOT use this device if a prescription has not been provided to you or if you do not understand the prescription. Instructions are printed on the top of the lid of your cold therapy device. Please refer to the lid for setup instructions.

WHAT DOES IT COST?

The unit is offered for **\$145.64** at the Time of Service. The device is yours at that point and may be used for future surgeries and/or injuries where a prescription for use is advised.

*Cold therapy devices are not covered by Medicare and are generally not covered by most insurances. The cost is \$145.64 at the Time of Service or we can submit to your insurance and you will be billed for whatever your insurance does not cover and may pay that portion in full or in payments once you receive your statement from DonJoy/DJO Global.

DIRECTION REMINDERS:

Always use a barrier between your skin and the cold therapy wrap when using the product to prevent over exposure to cold. Should you develop any type of abnormal rash that does not resolve and go away please discontinue use of the product.

THIS DEVICE CAN BE COLD ENOUGH TO CAUSE SERIOUS INJURY. READ AND UNDERSTAND ALL WARNINGS, LABELS AND INSTRUCTIONS FOR USE BEFORE USING THIS DEVICE. ADDITIONAL WARNINGS APPEAR IN THE INSTRUCTIONS FOR USE.

DONJOY® ULTRASLING™ ER IV



WHAT IS IT?

For protocols involving immobilization in external rotation, the UltraSling ER IV was designed to provide a comfortable proper fit—every time. Ideal for preferred positioning following:

- Capsular shifts
- Post-operative Bankart procedures
- Glenohumeral dislocations and subluxation
- Post-operative care after release of severe anterior capsule contracture
- Adhesive Capsulitis release and manipulation

MORE OPTIONS

- 15° and 30° cushions for solid, stable support.
- Cushions switch easily from right to left reducing inventory required.

UltraSling IV Product line leads the way in comfort and convenience:

- The three-point strapping system eliminates pressure on the neck
- Soft, breathable inner lining with mesh laminate helps improve ventilation
- Color-coded stress ball and label by size for quick identification and accurate dispensing
- Quick release buckles make it easy to take on and off for exercise/therapy

WHAT DOES IT COST?

We can submit to your insurance; you will be billed for whatever your insurance does not cover and may pay that portion in full or in payments once you receive your statement from DonJoy/DJO Global.

CLEANING INSTRUCTIONS:

Hand wash in cold water using mild soap, rinse thoroughly. Air dry. Note: If not rinsed thoroughly, residual soap may cause irritation and deteriorate material.

If you experience any pain, swelling, sensation changes, or any unusual reactions while using this product, consult your medical professional immediately. Never wear the support over an open wound.

Please contact your local DonJoy Territory Manager/DJO Representative with any questions.

**Anne Taylor (425) 691-7307
anne.taylor@djoglobal.com
*Please Contact 10 Days PRIOR to Surgery***

